



Higher Order Fibonacci Sequence and Series by Generalized Higher Order Variable Co-Efficient Difference Operator

Sandra Pinelas¹, Mohan B² and Britto Antony Xavier G^{3*}

¹Departamento de Cincias Exactas e Naturais Av. Conde Castro Guimares 2720-113, Amadora, Portugal.

^{2,3}Department of Mathematics, Sacred Heart College, Tirupattur, Vellore District- 635 601, TamilNadu, India.

Abstract

In this paper, we introduce generalized m^{th} order difference operator with variable co-efficient and its inverse by which we obtain higher Fibonacci sequence and its sum. Some theorems and interesting results on the sum of the terms of higher Fibonacci sequence with variable co-efficient are derived. Suitable examples are provided to illustrate our results.

Key words: Generalized difference operator, Variable co-efficient, Fibonacci sequence, Closed form solution, Fibonacci summation.

AMS classification: 39A70, 39A10, 47B39, 65J10, 65Q10.

1. Introduction

In 1984, Jerzy Popenda [5] introduced a particular type of difference operator on $u(k)$ as $\Delta_\alpha u(k) = u(k+1) - \alpha u(k)$. In 1989, Miller and Rose [8] introduced the discrete analogue of the Riemann-Liouville fractional derivative and its inverse $\Delta_h^{-\nu} f(t)$ ([1, 4]). In 2011, M.Maria Susai Manuel, et.al, [7] extended the operator Δ_α to generalized α -difference operator as $\Delta_{\alpha(\ell)} v(k) = v(k+\ell) - \alpha v(k)$ for the real valued function $v(k)$. In 2014, G.Britto Antony Xavier, et.al, [2] introduced q -difference operator as $\Delta_q v(k) = v(qk) - v(k)$, $q \in (0, \infty)$ and obtained finite series solution to the corresponding generalized q -difference equation $\Delta_q v(k) = u(k)$. With this background, in this paper, we obtain advanced Fibonacci sequence and its sum by introducing n^{th} -order difference operator with variable co-efficients.

^{3*}brittoshc@gmail.com

2. Higher order Fibonacci sequence and series by Generalized m-th order variable co-efficient difference equation

Fibonacci and Lucas numbers cover a wide range of interest in modern mathematics as they appear in the comprehensive works of Koshy [6] and Vajda [10]. The k -Fibonacci sequence introduced by Falcon and Plaza [3] depends only on one integer parameter k and is defined as

$$F_{k,0} = 0, \quad F_{k,1} = 1 \quad \text{and} \quad F_{k,n+1} = kF_{k,n} + F_{k,n-1}, \quad \text{where } n \geq 1, k \geq 1.$$

In particular, if $k = 2$, the Pell sequence is obtained as

$$P_0 = 0, \quad P_1 = 1 \quad \text{and} \quad P_{n+1} = 2P_n + P_{n-1} \quad \text{for } n \geq 1.$$

Here we introduce m^{th} -order generalized difference operator with variable co-efficients $\Delta_{\lambda_\alpha(\ell)} v(k) = v(k) - \sum_{i=1}^n \alpha_i k^{r_i} v(k - i\ell)$, $\lambda_\alpha(\ell) = (\alpha_1 k^{r_1}, \alpha_2 k^{r_2} \dots \alpha_m k^{r_m})$ which generates higher order Fibonacci sequence and its sum.

Definition 2.1 For $k \in [0, \infty)$, higher order Fibonacci sequence is defined as

$$F_0 = 1, \quad F_1 = \alpha_1 k^{r_1}, \quad F_n = \alpha_1 [k - (n-1)\ell]^{r_1} F_{n-1} + \alpha_2 [k - (n-2)\ell]^{r_2} F_{n-2}, \quad n \geq 2 \quad (1)$$

If $\alpha_1 = \alpha_2 = r_1 = r_2 = \ell = 1$ in (1), we have the well known Fibonacci sequence.

Example 2.2 (i) Taking $k = 7, \alpha_1 = 10, \alpha_2 = 7, r_1 = 3$ and $r_2 = 2$ in (1), we get a Fibonacci sequence $\{1, 490, 193207, 12173560, \dots\}$.

(ii) When $k = 9, \alpha_1 = 0.8, \alpha_2 = 0.3, r_1 = 2$ and $r_2 = 4$ in (1), we have a Fibonacci sequence $\{1, 583.2, 238903.02, 65566186.13, \dots\}$.

Similarly, one can obtain higher order Fibonacci sequences corresponding to each $\lambda_\alpha(\ell) = (\alpha_1 k^{r_1}, \alpha_1 k^{r_2} \dots \alpha_m k^{r_m}) \in \mathbb{R}^2$.

Definition 2.3 A generalized m^{th} order difference operator with variable co-efficients on $v(k)$, denoted as $\Delta_{\lambda_\alpha(\ell)} v(k)$, where $\lambda_\alpha(\ell) = (\alpha_1 k^{r_1}, \alpha_1 k^{r_2} \dots \alpha_m k^{r_m})$ is defined as

$$\Delta_{\lambda_\alpha(\ell)} v(k) = v(k) - \sum_{i=1}^m \alpha_i k^{r_i} v(k - i\ell), \quad k, \ell \in [0, \infty) \quad (2)$$

and its inverse is defined as below;

$$\text{if } \Delta_{\lambda_\alpha(\ell)} v(k) = u(k), \quad \text{then we write } v(k) = \overset{-1}{\Delta}_{\lambda_\alpha(\ell)} u(k). \quad (3)$$

Lemma 2.4 Let $v(k)$ be a functions of $k \in (-\infty, \infty)$. Then we obtain

$$\Delta_{\lambda_\alpha(\ell)}^{-1} a^{sk} \left[1 - \sum_{i=1}^m \frac{\alpha_i k^{r_i}}{a^{is\ell}} \right] = a^{sk}. \quad (4)$$

Proof: Taking $v(k) = a^{sk}$ in (2), we obtain $\Delta_{\lambda_\alpha(\ell)} a^{sk} = a^{sk} \left[1 - \sum_{i=1}^m \frac{\alpha_i k^{r_i}}{a^{is\ell}} \right]$. Now (4) follows from (3).

Corollary 2.5 If $m = 3$ in lemma (2.4), then we obtained

$$\Delta_{\lambda_\alpha(\ell)}^{-1} a^{sk} \left[1 - \sum_{i=1}^3 \frac{\alpha_i k^{r_i}}{a^{is\ell}} \right] = a^{sk}. \quad (5)$$

Proof: Taking $u(k) = a^{sk} \left[1 - \sum_{i=1}^3 \frac{\alpha_i k^{r_i}}{a^{is\ell}} \right]$ in (2), we have

$$\Delta_{\lambda_\alpha(\ell)} a^{sk} = a^{sk} \left[1 - \sum_{i=1}^3 \frac{\alpha_i k^{r_i}}{a^{is\ell}} \right].$$

Now (5) follows from (3).

Corollary 2.6 Let e^{-sk} be a function of $k \in (-\infty, \infty)$. Then we have

$$\Delta_{\lambda_\alpha(\ell)}^{-1} e^{-sk} \left[1 - \sum_{i=1}^m \alpha_i k^{r_i} e^{is\ell} \right] = e^{-sk}. \quad (6)$$

Proof: The proof follows by assuming $a = e^{-1}$ in (4).

Corollary 2.7 Let e^{-sk} be a function of $k \in (-\infty, \infty)$. Then we have

$$\Delta_{\lambda_\alpha(\ell)}^{-1} e^{-sk} \left[1 - \sum_{i=1}^3 \alpha_i k^{r_i} e^{is\ell} \right] = e^{-sk}. \quad (7)$$

Proof: The proof follows by assuming $m = 3$ in corollary (2.6).

Corollary 2.8 Let e^{sk} be a function of $k \in (-\infty, \infty)$, then we obtained

$$\Delta_{\lambda_\alpha(\ell)}^{-1} e^{sk} \left[1 - \sum_{i=1}^m \frac{\alpha_i k_i^r}{e^{is\ell}} \right] = e^{sk}. \quad (8)$$

Proof: The proof follows by taking $a = e^{sk}$ in lemma (2.4).

Corollary 2.9 Let e^{sk} be a function of $k \in (-\infty, \infty)$, then we obtained

$$\Delta_{\lambda_\alpha(\ell)}^{-1} e^{sk} \left[1 - \sum_{i=1}^3 \frac{\alpha_i k_i^r}{e^{is\ell}} \right] = e^{sk}. \quad (9)$$

Proof: The proof follows by taking $m = 3$ in corollary (2.8).

Corollary 2.10 Let $\log k$ be a function of $k > 2\ell$. Then we have

$$\Delta_{\lambda_\alpha(\ell)}^{-1} \left[\log k - \sum_{i=1}^m \alpha_i k_i^r \log(k - i\ell) \right] = \log k. \quad (10)$$

Proof: Taking $v(k) = \log k$ in (2), we obtained

$$\Delta_{\lambda_\alpha(\ell)} \log k = \log k - \sum_{i=1}^m \alpha_i k_i^r \log(k - i\ell).$$

Now (10) follows from (3).

Corollary 2.11 Let $\log k$ be a function of $k > 2\ell$. Then we have

$$\Delta_{\lambda_\alpha(\ell)}^{-1} \left[\log k - \sum_{i=1}^3 \alpha_i k_i^r \log(k - i\ell) \right] = \log k. \quad (11)$$

Proof: The proof follows by taking $m = 3$ in corollary (2.10).

Theorem 2.12 If $v(k) = \Delta_{\lambda_\alpha(\ell)}^{-1} u(k)$, $F_0 = 1$, $F_1 = F_0 \alpha_1 k^{r_1}$ and

$$F_{n+1} = \sum_{i=0}^n F_{n-i} \alpha_{i+1} [k - (n-i)\ell]^{r_{i+1}}, \text{ then } \sum_{i=0}^n F_i u(k - i\ell) = v(k) - F_{n+1} v(k - (n+1)\ell) -$$

$$\sum_{i=0}^n F_{n-i} \alpha_{i+2} [k - (n-i)\ell]^{r_{i+2}} v(k - (n+2)\ell) -$$

$$\sum_{i=0}^{n-1} F_{n-i} \alpha_{i+3} [k - (n-i)\ell]^{r_{i+3}} v(k - (n+3)\ell) + \dots + F_n \alpha_m (k - n\ell)^{r_m} v(k - (n+m)\ell). \quad (12)$$

Proof: From (2) and (3), we arrive

$$v(k) = u(k) + \alpha_1 k^{r_1} v(k - \ell) + \alpha_2 k^{r_2} v(k - 2\ell) + \dots + \alpha_m k^{r_m} v(k - m\ell). \quad (13)$$

Replacing k by $k - \ell$ and then substituting the value of $v(k - \ell)$ in (13), we get

$$v(k) = u(k) + F_1 u(k - \ell) + [F_1 \alpha_1 (k - \ell)^{r_1} + \alpha_2 k^{r_2}] v(k - 2\ell) + \dots +$$

$$[F_1 \alpha_{m-1} (k - \ell)^{r_{m-1}} + \alpha_m k^{r_m}] v(k - m\ell) + F_1 \alpha_m (k - \ell)^{r_m} v(k - (m + 1)\ell) \quad (14)$$

which gives

$$v(k) = u(k) + F_1 u(k - \ell) + F_2 v(k - 2\ell) + [F_1 \alpha_2 (k - \ell)^{r_2} + \alpha_3 k^{r_3}] v(k - 3\ell) + \dots +$$

$$[F_1 \alpha_{m-1} (k - \ell)^{r_{m-1}} + \alpha_m k^{r_m}] v(k - m\ell) + F_1 \alpha_m (k - \ell)^{r_m} v(k - (m + 1)\ell), \quad (15)$$

where F_0 , F_1 and F_2 are given in (1).

Replacing k by $k - 2\ell$ in (13) and then substituting $v(k - 2\ell)$ in (15), we obtain

$$v(k) = \sum_{i=0}^3 F_i u(k - i\ell) + \dots + F_2 \alpha_m (k - 2\ell)^{r_m} v(k - (m + 2)\ell),$$

where F_3 is given in (1). Repeating this process again and again, we get (12).

Corollary 2.13 If $v(k) = \Delta_{\lambda_\alpha(\ell)}^{-1} u(k)$, $F_0 = 1$, $F_1 = F_0 \alpha_1 k^{r_1}$ and

$$F_{n+1} = \sum_{i=0}^n F_{n-i} \alpha_{i+1} [k - (n-i)\ell]^{r_{i+1}}, \text{ then } \sum_{i=0}^n F_i u(k - i\ell) = v(k) - F_{n+1} v(k - (n+1)\ell) -$$

$$\sum_{i=0}^1 F_{n-i} \alpha_{i+2} [k - (n-i)\ell]^{r_{i+2}} v(k - (n+2)\ell) + F_n \alpha_3 (k - n\ell)^{r_3} v(k - (n+3)\ell). \quad (16)$$

Proof: The proof follows by taking $m = 3$ in Theorem (2.12).

Corollary 2.14 If $v(k)$ is a closed form solution of the m^{th} order generalized difference equation

$$\Delta_{\lambda_\alpha(\ell)} v(k) = a^{sk} \left[1 - \frac{\alpha_1 k^{r_1}}{a^{s\ell}} - \frac{\alpha_2 k^{r_2}}{a^{2s\ell}} - \frac{\alpha_3 k^{r_3}}{a^{3s\ell}} \right],$$

then we obtain

$$\begin{aligned}
 a^{sk} & \left[1 - \frac{F_{n+1}}{a^{s(n+1)\ell}} - \sum_{i=0}^1 \frac{F_{n-i}\alpha_{i+2}[k - (n-i)\ell]^{r_{i+2}}}{a^{s(n+2)\ell}} - \frac{F_n\alpha_3(k-n\ell)^{r_3}}{a^{s(n+3)\ell}} \right] \\
 & = \sum_{i=0}^n F_i a^{s(k-i\ell)} \left[1 - \frac{\alpha_1(k-i\ell)^{r_1}}{a^{s\ell}} - \frac{\alpha_2(k-i\ell)^{r_2}}{a^{2s\ell}} - \frac{\alpha_3(k-i\ell)^{r_3}}{a^{3s\ell}} \right]. \quad (17)
 \end{aligned}$$

Proof: The proof of (17) follows by taking $v(k) = a^{sk}$ and applying (4) in (16).

The following example is an verification of corollary 2.14.

Example 2.15 Taking $k = 9, \ell = 0.3, n = 1, a = 5, \alpha_1 = 0.2, \alpha_2 = 0.3, \alpha_3 = 0.4, r_1 = 1$ and $r_2 = 3, r_3 = 4$ in (17), we get

$$5^9 - F_2 5^{-2} - 3F_2 5^{-5} = \sum_{i=0}^1 F_i 5^{(9-0.3i)} \left[1 - \frac{2(7-3i)^1}{5^3} - \frac{3(7-3i)^2}{5^6} \right] = 78077.15136,$$

where $F_0 = 1, F_1 = 14, F_2 = 259, F_3 = 1190$.

Corollary 2.16 Let e^{-sk} be a function of $k \in (-\infty, \infty)$. Then

$$\begin{aligned}
 e^{-sk} & \left[1 - F_{n+1} e^{s(n+1)\ell} - \sum_{i=0}^1 F_{n-i}\alpha_{i+2}[k - (n-i)\ell]^{r_{i+2}} e^{s(n+2)\ell} + F_n\alpha_3(k-n\ell)^{r_3} e^{s(n+3)\ell} \right] \\
 & = \sum_{i=0}^n F_i e^{-s(k-i\ell)} \left[1 - \alpha_1(k-i\ell)^{r_1} e^{s\ell} - \alpha_2(k-i\ell)^{r_2} e^{2s\ell} - \alpha_3(k-i\ell)^{r_3} e^{3s\ell} \right]. \quad (18)
 \end{aligned}$$

Proof: Taking $v(k) = e^{-sk}$ and applying (6) in (4), we get (18).

Example 2.17 Taking $k = 9, \ell = 1, n = 3, \alpha_1 = 0.8, \alpha_2 = 0.3, r = 3$ and $s = 2$ in (18), then we obtained

$$e^{-9} - F_4 e^5 - (0.3)6^2 F_3 e^{-4} = \sum_{i=0}^3 F_i e^{-(9-i)} \left[1 - (0.8)(9-i)^3 e - (0.3)(9-i)^2 e^2 \right] = -89333078.94$$

where $F_0 = 1, F_1 = 583.2, F_2 = 238903.02, F_3 = 65566186.13$ and $F_4 = 11333348840$.

Theorem 2.18 Let $t \in \mathbb{N}(0)$. Then a closed form solution of the generalized m^{th} order difference equation $v(k) - \sum_{i=1}^m \alpha_i k^{r_i} v(k-i\ell) = \left[k^t - \sum_{i=1}^m \alpha_i k^{r_i} (k-i\ell)^t \right]$ is

$$\Delta_{\lambda_\alpha(\ell)}^{-1} \left[k^t - \sum_{i=1}^m \alpha_i k^{r_i} (k-i\ell)^t \right] = k^t \quad (19)$$

Proof: Taking $v(k) = k^t$ in (2) and using (3), we get (19).

Corollary 2.19 If $v(k) = \Delta_{\lambda\alpha(\ell)}^{-1} \left[k^t - \sum_{p=1}^m \alpha_p k^{r_p} (k - p\ell)^t \right]$ is the closed form solution given in (19), then

$$v(k) - F_{n+1}(k - (n + 1)\ell)^t - \sum_{i=0}^n F_{n-i}\alpha_{i+2}[k - (n - i)\ell]^{r_{i+2}}(k - (n + 2)\ell)^t + \dots + F_n\alpha_m(k - n\ell)^{r_m}(k - (n + m)\ell)^t = \sum_{i=0}^n F_i \left[(k - i\ell)^t - \sum_{p=1}^m \alpha_p (k - p\ell)^{r_p} [k - (i + p)\ell]^t \right]. \quad (20)$$

Proof: Taking $u(k) = k^t - \sum_{p=1}^m \alpha_p k^{r_p} (k - p\ell)^t$ in Theorem (2.12), we have (20).

Example 2.20 Let $k = 7$, $\ell = 2$, $n = 3$, $t = 2$, $r = 3$, $s = 4$ $\alpha_1 = 5$, $\alpha_2 = 3$ in Corollary (2.19). Then

$$\sum_{i=0}^3 F_i u(7 - 2i) = v(7) - F_4 v(-1) - \alpha_2 F_3 v(-3) = -5, 026, 731, 585.$$

where $u(k) = k^t - \alpha_1 k^r (k - \ell)^t - \alpha_2 k^s (k - 2\ell)^t$, $F_0 = 1$, $F_1 = 1715$, $F_2 = 1, 079, 078$, $F_3 = 148, 891, 115$ and $F_4 = 1, 006, 671, 529$.

Corollary 2.21 If $v(k)$ is a closed form solution of m^{th} order difference equation with variable co-efficients

$$v(k) - \sum_{i=1}^m \alpha_i k^{r_i} v(k - i\ell) = k^t a^{sk} - \sum_{i=1}^m \left[\alpha_i k^{r_i} (k - i\ell)^t a^{s(k-i\ell)} \right], \text{ then we have}$$

$$k^t a^{sk} - F_{n+1}(k - (n + 1)\ell)^t a^{s(k-(n+1)\ell)} - \sum_{i=0}^n F_{n-i}\alpha_{i+2}[k - (n - i)\ell]^{r_{i+2}} \times (k - (n + 2)\ell)^t a^{s(k-(n+2)\ell)} + \dots + F_n\alpha_m(k - n\ell)^{r_m}(k - (n + m)\ell)^t a^{s(k-(n+m)\ell)}$$

$$= \sum_{i=0}^n F_i \left[(k - i\ell)^t - \sum_{p=1}^m \alpha_p (k - p\ell)^{r_p} [k - (i + p)\ell]^t a^{s[k-(i+p)\ell]} \right]. \quad (21)$$

Proof: Taking $u(k) = k^t a^{sk} - \sum_{i=1}^m \left[\alpha_i k^{r_i} (k - i\ell)^t a^{s(k-i\ell)} \right]$ in Theorem (2.12) and using (4), we get (21).

Corollary 2.22 A closed form solution of generalized third order difference equation

$$\Delta_{\lambda\alpha(\ell)} v(k) = k^2 a^{sk} - \sum_{i=1}^3 \left[\alpha_i k^{r_i} (k - i\ell)^2 a^{s(k-i\ell)} \right] \text{ is } k^2 a^k \text{ and hence we have}$$

$$\begin{aligned}
 & k^2 a^{sk} - F_{n+1}(k - (n + 1)\ell)^2 a^{s(k-(n+1)\ell)} - \sum_{i=0}^1 F_{n-i} \alpha_{i+2} [k - (n - i)\ell]^{r_{i+2}} \\
 & (k - (n + 2)\ell)^2 a^{s(k-(n+2)\ell)} + F_n \alpha_3 (k - n\ell)^{r_3} (k - (n + 3)\ell)^2 a^{s(k-(n+3)\ell)} \\
 & = \sum_{i=0}^n F_i \left[(k - i\ell)^2 a^{s(k-i\ell)} - \sum_{p=1}^3 \alpha_p (k - i\ell)^{r_p} [k - (p + i)\ell]^2 a^{s(k-(p+i)\ell)} \right]. \quad (22)
 \end{aligned}$$

Proof: The proof follows by taking $m = 3$ and $t = 2$ in Corollary (2.21).

Example 2.23 Let $k = 5$, $\ell = 2$, $a = 3$, $n = 4$, $\alpha_1 = 0.02$, $\alpha_2 = 0.03$, $r = 3$, $s = 2$ in Corollary (2.22). Then we obtain

$$\begin{aligned}
 v(5) - F_5 v(-5) - (0.03) F_4 v(-7) &= \sum_{i=0}^3 F_i [(5 - 2i)^3 3^{k-2i} - (0.02)(5 - 2i)^3 \times \\
 [5 - (i + 1)2]^3 3^{5-(i+1)2} - (.03)(5 - 2i)^2 [5 - 2(i + 2)]^3 3^{5-2(i+2)}] &= 24, 611, 856.47,
 \end{aligned}$$

where $F_0 = 1$, $F_1 = 2.5$, $F_2 = 2.1$, $F_3 = 0.717$, $F_4 = 0.04866$ and $F_5 = 0.0477864$.

Corollary 2.24 A closed form solution of the second order difference equation

$$\begin{aligned}
 v(k) - \sum_{i=0}^3 \alpha_i k^{r_i} v(k - i\ell) &= k^t e^{-sk} - \sum_{p=0}^3 \alpha_p k^{r_p} (k - p\ell)^t e^{-s(k-p\ell)} \text{ is given by} \\
 & k^t e^{-sk} - F_{n+1}(k - (n + 1)\ell)^t e^{-s(k-(n+1)\ell)} - \sum_{i=0}^1 F_{n-i} \alpha_{i+2} [k - (n - i)\ell]^{r_{i+2}} \times \\
 & (k - (n + 2)\ell)^t e^{-s(k-(n+2)\ell)} + F_n \alpha_3 (k - n\ell)^{r_3} (k - (n + 3)\ell)^t e^{-s(k-(n+3)\ell)} \\
 & = \sum_{i=0}^n F_i e^{-s(k-i\ell)} \left[(k - i\ell)^t - \sum_{p=1}^3 \alpha_p (k - (p + i)\ell)^{r_p} [k - (p + i)\ell]^t e^{sp\ell} \right]. \quad (23)
 \end{aligned}$$

Proof: Taking $a = e^{-1}$ in (21), we get (23).

Corollary 2.25 If $v(k) = \underset{\lambda_\alpha(\ell)}{\Delta}^{-1} \left[k e^{-sk} - \sum_{p=1}^m \alpha_p k^p (k - p\ell) e^{-s(k-\ell)} \right]$ is the closed form solution given in (23), then

$$\begin{aligned}
 & k e^{-sk} - F_{n+1}(k - (n + 1)\ell) e^{-s(k-(n+1)\ell)} - \sum_{i=0}^1 F_{n-i} \alpha_{i+2} [k - (n - i)\ell]^{r_{i+2}} \times \\
 & (k - (n + 2)\ell) e^{-s(k-(n+2)\ell)} + F_n \alpha_3 (k - n\ell)^{r_3} (k - (n + 3)\ell) e^{-s(k-(n+3)\ell)} \\
 & = \sum_{i=0}^n F_i e^{-s(k-i\ell)} \left[(k - i\ell) - \sum_{p=1}^3 \alpha_p (k - (p + i)\ell)^{r_p} [k - (p + i)\ell] e^{sp\ell} \right]. \quad (24)
 \end{aligned}$$

Proof: The proof follows by taking $t = 1$ in Corollary (2.24).

Theorem 2.26 Let $v(k)$ be a solution of the n^{th} -order difference equation with variable co-efficients

$$v(k) - \sum_{i=0}^m \alpha_i k^{r_i} v(k - i\ell) = k^{(t)} a^{sk} - \sum_{p=1}^m \alpha_p k^{r_p} (k - p\ell)^{(t)} a^{s(k-p\ell)},$$

then we have

$$\begin{aligned} & k^{(t)} a^{sk} - F_{n+1}(k - [n + 1]\ell)^{(t)} a^{s(k-[n+1]\ell)} - \sum_{i=0}^n F_{n-i} \alpha_{i+2} [k - (n - i)\ell]^{r_{i+2}} \times \\ & (k - [n + 2]\ell)^{(t)} a^{s(k-[n+2]\ell)} - \sum_{i=0}^{n-1} F_{n-i} \alpha_{i+3} [k - (n - i)\ell]^{r_{i+3}} (k - [n + 3]\ell)^{(t)} a^{s(k-[n+3]\ell)} \\ & + \dots + F_n \alpha_m (k - n\ell)^{r_m} (k - [n + m]\ell)^{(t)} a^{s(k-[n+m]\ell)} \\ & = \sum_{i=0}^n F_i a^{s(k-i\ell)} \left[(k - i\ell)^{(t)} - \sum_{p=1}^m \alpha_p (k - i\ell)^{r_p} (k - (i + p)\ell)^{(t)} a^{-ps\ell} \right] \end{aligned} \quad (25)$$

Proof: Taking $v(k) = k^{(t)} a^{sk}$ in Theorem (2.12) and using (4), we get (25).

Corollary 2.27 If $v(k)$ is the closed form solution given of (25), then

$$\begin{aligned} & k^{(2)} a^{sk} - F_{n+1}(k - [n + 1]\ell)^{(2)} a^{s(k-[n+1]\ell)} - \sum_{i=0}^1 F_{n-i} \alpha_{i+2} [k - (n - i)\ell]^{r_{i+2}} \times \\ & (k - (n + 2)\ell)^{(2)} a^{-s(k-(n+2)\ell)} - F_n \alpha_3 (k - n\ell)^{r_3} (k - (n + 3)\ell) e^{-s(k-(n+3)\ell)} \\ & = \sum_{i=0}^n F_i a^{s(k-i\ell)} \left[(k - i\ell)^{(2)} - \sum_{p=1}^3 \alpha_p (k - i\ell)^{r_p} [k - (i + p)\ell]^{(2)} a^{-ps\ell} \right]. \end{aligned} \quad (26)$$

Proof: The proof follows by taking $m = 3$ and $t = 2$ in Theorem (2.26).

Example 2.28 Let $k = 7$, $\ell = 2$, $a = 3$, $n = 2$, $\alpha_1 = 0.04$, $\alpha_2 = 0.06$, $r = 4$, $s = 3$ in Corollary (2.27). Then we obtain

$$\begin{aligned} & v(7) - F_3 v(1) - (0.06) 3^3 F_2 v(-1) = \sum_{i=0}^2 F_i [(7 - 2i)^{(2)} 3^{k-2i} - (0.06)(7 - 2i)^3 \\ & [7 - 2(i + 1)]^{(2)} 3^{7-2(i+1)} - (.06)(7 - 2i)^{(2)} [7 - 2(i + 2)]^3 3^{7-2(i+2)}] = 84008.0808, \end{aligned}$$

where $F_0 = 1$, $F_1 = 96.04$, $F_2 = 2421.58$, $F_3 = 8566.2192$.

Corollary 2.29 Let $v(k)$ be a solution of n^{th} order difference equation with variable

$$v(k) - \sum_{i=0}^m \alpha_i k^{r_i} v(k - i\ell) = e^{-sk} \left[k^{(2)} - \sum_{p=1}^m \alpha_p k^{r_p} (k - p\ell)^{(2)} e^{ps\ell} \right].$$

Then we have

$$\begin{aligned}
 & k^{(2)}e^{-sk} - F_{n+1}(k - [n + 1]\ell)^{(2)}e^{-s(k-[n+1]\ell)} - \sum_{i=0}^1 F_{n-i}\alpha_{i+2}[k - (n - i)\ell]^{r_{i+2}} \times \\
 & (k - (n + 2)\ell)^{(2)}e^{-s(k-(n+2)\ell)} - F_n\alpha_3(k - n\ell)^{r_3}(k - (n + 3)\ell)^{(2)}e^{-s(k-(n+3)\ell)} \\
 & = \sum_{i=0}^n F_i e^{-(k-i\ell)} \left[(k - i\ell)^{(2)} - \sum_{p=1}^m \alpha_p (k - i\ell)^{r_p} [k - (i + p)\ell]^{(2)} e^{ps\ell} \right]. \quad (27)
 \end{aligned}$$

Proof: Taking $a = e^{-1}$ in (2.27), we get (27).

Example 2.30 Let $k = 6, \ell = 0.21, n = 2, a = 0.2, \alpha_1 = 2, \alpha_2 = 0.3, r = 3, s = 2$ in Corollary (2.29). Then we obtain

$$\begin{aligned}
 v(6) - F_3 v(5.37) - (0.3)(5.58)^2 F_2 v(5.16) &= \sum_{i=0}^3 F_i [(6 - (0.21)i)^{(3)}(0.2)^{k-(0.21)i} - \\
 & (2)(6 - (0.21)i)^3 [6 - (0.21)(i + 1)]^{(3)}(0.2)^{6-(0.21)(i+1)} - (.3)(6 - (0.21)i)^2 \\
 & [6 - (0.21)(i + 2)]^{(3)}3^{6-(0.21)(i+2)}] = -7, 539, 276.7060093,
 \end{aligned}$$

where $F_0 = 1, F_1 = 432, F_2 = 167717.1217$ and $F_3 = 8746152.49$.

3. Conclusion

We obtained summation formula to Higher order Fibonacci sequence by introducing generalized m^{th} order difference operator with variable co-efficients and have derived certain results on closed and summation form solution of generalized m^{th} order difference equation with variable co-efficients which will be used to our further research.

References

- [1] Bastos NRO, Ferreira RAC and Torres DFM, Discrete-Time Fractional Variational Problems, Signal Processing, 91(3), 2011, 513-524.
- [2] Britto Antony Xavier G, Gerly TG and Nasira Begum H, Finite Series of Polynomials and Polynomial Factorials arising from Generalized q-Difference operator, Far East Journal of Mathematical Sciences, 94(1), 2014, 47-63.
- [3] Falcon S and Plaza A, On the Fibonacci k -numbers, Chaos Solitons and Fractals, 32(5), 1615-1624, (2007).
- [4] Ferreira RAC and Torres DFM, Fractional h-difference equations arising from the calculus of variations, Applicable Analysis and Discrete Mathematics, 5(1), 2011, 110-121.

- [5] Jerzy Popenda and Blazej Szmanda, On the Oscillation of Solutions of Certain Difference Equations, *Demonstratio Mathematica*, XVII(1), 1984, 153 - 164.
- [6] Koshy T, *Fibonacci and Lucas Numbers with Applications*, Wiley-Inter science, New York, NY, USA, (2001).
- [7] Maria Susai Manuel M, Chandrasekar V and Britto Antony Xavier G, Solutions and Applications of Certain Class of α -Difference Equations, *International Journal of Applied Mathematics*, 24(6), 2011, 943-954.
- [8] Miller KS and Ross B, *Fractional Difference Calculus in Univalent Functions*, Horwood, Chichester, UK, 139-152, (1989).
- [9] Susai Manuel M, Britto Antony Xavier G, Chandrasekar V and Pugalarasu R, Theory and application of the Generalized Difference Operator of the n^{th} kind(Part I), *Demonstratio Mathematica*, 45(1), 2012, 95-106.
- [10] Vajda S, *Fibonacci and Lucas Numbers, and the Golden Section*, Ellis Horwood, Chichester, UK, (1989).