



Forced Oscillation of Solutions of Fractional Neutral Nonlinear Partial Differential Equations

V.Sadhasivam^{1*}, Kavitha J² and Nagajothi N³

^{1,2,3}Department of Mathematics, Thiruvalluvar Government Arts College,
Rasipuram - 637 401, Namakkal Dt., Tamil Nadu, S.India.

Abstract

In this article, we investigate the oscillation of fractional order neutral partial differential equations of the form $D_{+,t}^{1+\alpha} [u(x,t) + \lambda(t)u(x,t-\sigma)] + r(t)D_{+,t}^{\alpha} u(x,t) = a(t)\Delta u(x,t) + \sum_{j=1}^l a_j(t)\Delta u(x,t-\tau_j) - p(x,t)F\left(\int_0^t (t-s)^{-\alpha} u(x,s)ds\right) + f(x,t)$, $(x,t) \in \Omega \times R_+ = G$. Using the generalized Riccati technique and integral averaging method, new oscillation criteria are established.

AMS Classification: 34K37, 35B05, 35R11.

1. Introduction

Neutral differential equations are functional differential equation in which the highest order derivative of the unknown function appear both with and without deviations. The neutral differential equations arise in many areas of applied mathematics. During the last two decades there has been a lot of interest towards the study of qualitative theory of neutral differential equations. A good guide concerning the literature for ordinary neutral functional differential equation [14,29] and the references cited therein. The neutral delay differential equations arise in modeling of the networks containing loss less transmission lines (as in high-speed switching circuits), second order neutral delay differential equations appear in the study of vibrating masses attached to an elastic bar, as the Euler equation in some variational problems, theory of automatic control and in neuro mechanical systems in which inertia plays an important role. See [16,34].

On the other hand, for partial neutral functional differential equations we refer to the reader to [15,35] and the references cited therein. The investigation of the problem of oscillation and nonoscillation of neutral delay differential equations has been initiated by [36]. Recently, researchers have established some oscillation results for partial functional equations we refer the reader to [3,11,22] for parabolic equation and to [2,4-6,19] for hyperbolic equations.

^{1*}ovsadha@gmail.com, ²kaviakshita@gmail.com

In the last few years, the problem of oscillation of solutions of fractional order differential equation have received a great deal of attention. we refer in particular to the papers [7,10,12,13]. For general back round on fractional differential equations (see the monographs [1, 8, 9, 20, 21, 23, 29, 30, 31,37]). However, it seems that very little is known regarding the oscillatory behavior of fractional order partial differential equations [18,24-27,32,33] and the references cited therein.

In [33], Wei Nian Li investigated the oscillation properties for solutions of a kind of partial fractional differential equations with damping term of the form

$$D_{+,t}^{1+\alpha} u(x, t) + p(t)D_{+,t}^{\alpha} (u(x, t)) = a(t)\Delta u(x, t) + \sum_{i=1}^m a_i(t)\Delta u(x, t - \tau_i) - q(x, t) \left(\int_0^t (t - \xi)^{-\alpha} u(x, \xi) d\xi \right), \quad (x, t) \in G = \Omega \times R_+.$$

It seems that there has been no attempt made on $(1 + \alpha)^{th}$ order fractional neutral partial differential equation. Motivated by this, we study the following equation (1.1). In this paper, we obtain some new oscillation criteria for fractional order neutral partial differential equations with damping term of the form

$$D_{+,t}^{1+\alpha} [u(x, t) + \lambda(t)u(x, t - \sigma)] + r(t)D_{+,t}^{\alpha} u(x, t) = a(t)\Delta u(x, t) + \sum_{j=1}^l a_j(t)\Delta u(x, t - \tau_j) - p(x, t)F \left(\int_0^t (t - s)^{-\alpha} u(x, s) ds \right) + f(x, t), \quad (x, t) \in \Omega \times R_+ = G, \quad (1)$$

Ω is a bounded domain in R^n with a piecewise smooth boundary $\partial\Omega$, $\alpha \in (0, 1)$ is a constant, $D_{+,t}^{\alpha}$ is the Riemann-Liouville fractional derivative of order α of u with respect to t and $\Delta u(x, t) = \sum_{r=1}^n \frac{\partial^2 u(x, t)}{\partial x_r^2}$ is the Laplacian operator in the Euclidean n - space R^n .

Equation (1.1) is associated with the boundary conditions, namely

$$\frac{\partial u(x, t)}{\partial \gamma} + \psi(x, t)u(x, t) = 0, \quad (x, t) \in \partial\Omega \times R_+, \quad (2)$$

where γ is the unit outward drawn normal vector to $\partial\Omega$ and $\psi(x, t)$ is nonnegative continuous function on $\partial\Omega \times R_+$ and

$$u(x, t) = 0, \quad (x, t) \in \partial\Omega \times R_+. \quad (3)$$

We assume the following conditions throughout this paper without mentioning that

- (A₁) $\lambda \in C^{1+\alpha}([0, \infty); [0, \infty))$, $0 \leq \lambda < 1$ and σ is a nonnegative constant;
- (A₂) $r \in C([0, \infty); [0, \infty))$, $a \in C([0, \infty); [0, \infty))$, $a_j \in C([0, \infty); [0, \infty))$ and τ_j are

^{1*}ovsadha@gmail.com, ²kaviakshita@gmail.com

- nonnegative constant, $j = 1, 2, 3, \dots, l$;
 (A₃) $p \in C(\bar{G}; R_+)$ and $p(t) = \min_{x \in \bar{\Omega}} p(x, t)$;
 (A₄) $F \in C(R; R)$ is convex in $[0, \infty)$, and $uF(u) > 0$ for $u \neq 0$ and there exists a positive constants ρ such that $\frac{F(u)}{u} \geq \rho$ for $u \neq 0$;
 (A₅) $f \in C(\bar{G}; R)$ such that $\int_{\Omega} f(x, t) dx \leq 0$.

By a solution of the problem (1.1),(1.2) (or (1.1),(1.3)), we mean a function $u(x, t) \in C^{1+\alpha}(G) \cap C^{\alpha}(\bar{G})$ which satisfies (1.1) on G and the associated boundary condition (1.2)(or (1.3)).

The solution $u(x,t)$ of (1.1), (1.2) or (1.1), (1.3) is said to be oscillatory in the domain G if for any positive number θ there exists a point $(x_0, t_0) \in \Omega \times [\theta, \infty)$ such that $u(x_0, t_0) = 0$ holds.

Our main aim of this paper is to establish new oscillation criteria for (1.1), (1.2) and (1.1), (1.3) by using generalized Riccati technique method.

2. Preliminaries

In this section, we give the definitions of fractional derivatives and integrals which are useful throughout this paper, there are several kinds of definitions of fractional derivatives and integrals. In this paper, we use the Riemann-Liouville left sided definition on the half axis R_+ .

Definition 2.1 [28] The Riemann-Liouville fractional partial derivative of order $0 < \alpha < 1$ with respect to t of a function $u(x, t)$ is given by

$$(D_{+,t}^{\alpha} u)(x, t) := \frac{\partial}{\partial t} \frac{1}{\Gamma(1-\alpha)} \int_0^t (t-\nu)^{-\alpha} u(x, \nu) d\nu \quad (4)$$

provided the right hand side is pointwise defined on R_+ , where Γ is the gamma function.

Definition 2.2 [28] The Riemann-Liouville fractional integral of order $\alpha > 0$ of a function $y : R_+ \rightarrow R$ on the half-axis R_+ is given by

$$(I_+^{\alpha} y)(t) := \frac{1}{\Gamma(\alpha)} \int_0^t (t-\nu)^{\alpha-1} y(\nu) d\nu \quad \text{for } t > 0 \quad (5)$$

provided the right hand side is pointwise defined on R_+ .

Definition 2.3 [28] The Riemann-Liouville fractional derivative of order $\alpha > 0$ of a

function $y : R_+ \rightarrow R$ on the half-axis R_+ is given by

$$(D_+^\alpha y)(t) := \frac{d^{[\alpha]}}{dt^{[\alpha]}} \left(I_+^{[\alpha]-\alpha} y \right) (t) \quad \text{for } t > 0 \quad (6)$$

provided the right hand side is pointwise defined on R_+ , where $[\alpha]$ is the ceiling function of α .

We need the following lemmas in proving our main results.

Lemma 2.4 [15] Let

$$K(t) := \int_0^t (t - \nu)^{-\alpha} y(\nu) d\nu \quad \text{for } \alpha \in (0, 1) \quad \text{and } t > 0. \quad (7)$$

Then $K'(t) = \Gamma(1 - \alpha)(D_+^\alpha y)(t)$.

Lemma 2.5 [28] Let $\alpha \geq 0, m \in \mathbb{N}$ and $D = \frac{d}{dt}$. If the fractional derivatives $(D_+^\alpha y)(t)$ and $(D_+^{\alpha+m} y)(t)$ exists

$$D^m((D_+^\alpha y)(t)) = (D_+^{\alpha+m} y)(t). \quad (8)$$

Lemma 2.6 [27] If X and Y are nonnegative, then

$$mXY^{m-1} - X^m \leq (m - 1)Y^m. \quad (9)$$

3. Main Results

In this section, we establish the oscillation of the problem (1.1), (1.2). We start with the following theorem.

Theorem 3.1 Suppose that the conditions $(A_1) - (A_5)$ hold. Assume that for some $t_0 > 0$ and $F'(v)$ exists such that $F'(v) \geq \mu$ for some $\mu > 0$ and for all $v \neq 0$,

$$\int_{t_0}^{\infty} \exp \left(\int_{t_0}^t r(s) ds \right) dt = \infty, \quad (10)$$

$$\limsup_{t \rightarrow \infty} \int_{t_1}^t \left[v(s)p(s) - \frac{(r(s))^2 v(s)}{4\mu\Gamma(1 - \alpha)} \right] ds = \infty, \quad (11)$$

^{1*}ovsadha@gmail.com, ²kaviakshita@gmail.com

where

$$v(t) = \exp \left(- \int_{t_0}^t r(s) ds \right), t \geq t_0. \quad (12)$$

Then every solution of the problem (1.1), (1.2) is oscillatory in G.

Proof: Let us prove this theorem by the method of contradiction. Assume that there is a non-oscillatory solution $u(x, t)$ to the problem (1.1), (1.2) which has no zero in $\Omega \times [t_0, \infty)$ for some $t_0 > 0$. Without loss of generality we may suppose that $u(x, t) > 0, u(x, t - \tau_j) > 0$ and $u(x, t - \sigma) > 0$ in $\Omega \times [t_1, \infty)$ for $t_1 \geq t_0, j = 1, 2, 3, \dots, l$. Integrating (1.1) over the domain Ω , we get

$$\begin{aligned} & \int_{\Omega} D_{+,t}^{1+\alpha} [u(x, t) + \lambda(t)u(x, t - \sigma)] dx + r(t) \int_{\Omega} D_{+,t}^{\alpha} u(x, t) dx = a(t) \int_{\Omega} \Delta u(x, t) dx \\ & + \sum_{j=1}^l a_j(t) \int_{\Omega} \Delta u(x, t - \tau_j) dx - \int_{\Omega} p(x, t) F \left(\int_0^t (t-s)^{-\alpha} u(x, s) ds \right) dx \\ & + \int_{\Omega} f(x, t) dx, \quad t \geq t_1. \end{aligned} \quad (13)$$

Using Green's formula and (1.2), it follows

$$\int_{\Omega} \Delta u(x, t) dx = \int_{\partial\Omega} \frac{\partial u(x, t)}{\partial \gamma} dS = - \int_{\partial\Omega} \psi(x, t) u(x, t) dS \leq 0, \quad t \geq t_1 \quad (14)$$

and

$$\begin{aligned} \int_{\Omega} \Delta u(x, t - \tau_j) dx &= \int_{\partial\Omega} \frac{\partial u(x, t - \tau_j)}{\partial \gamma} dS \\ &= - \int_{\partial\Omega} \psi(x, t - \tau_j) u(x, t - \tau_j) dS \leq 0, \quad t \geq t_1, \quad j = 1, 2, 3, \dots, l. \end{aligned} \quad (15)$$

where dS is the surface element on $\partial\Omega$.

$$\text{Let } V(t) = \int_{\Omega} u(x, t) dx. \quad (16)$$

By using Jensen's inequality and (A_3) , we obtain

$$\begin{aligned} \int_{\Omega} p(x, t) F \left(\int_0^t (t-s)^{-\alpha} u(x, s) ds \right) dx &\geq p(t) F \left(\int_{\Omega} \left(\int_0^t (t-s)^{-\alpha} u(x, s) ds \right) dx \right) \\ &\geq p(t) F \left(\int_0^t (t-s)^{-\alpha} \left(\int_{\Omega} u(x, s) dx \right) ds \right) \\ &\geq p(t) F (K(t)), \quad t \geq t_1. \end{aligned} \quad (17)$$

By (A_5) ,

$$\int_{\Omega} f(x, t) dx \leq 0. \quad (18)$$

Combining (3.5) – (3.9), we have

$$D_+^{1+\alpha} [V(t) + \lambda(t)V(t - \sigma)] + r(t)D_+^{\alpha}V(t) \leq -p(t)F (K(t)), \quad t \geq t_1.$$

Let

$$z(t) = V(t) + \lambda(t)V(t - \sigma). \quad (19)$$

Then the above inequality becomes,

$$D_+^{1+\alpha} [z(t)] + r(t)D_+^{\alpha}V(t) + p(t)F (K(t)) \leq 0, \quad t \geq t_1. \quad (20)$$

Using Lemma 2.2 and (3.11), we have

$$\begin{aligned} [(D_+^{\alpha}z(t)) v(t)]' &= D_+^{1+\alpha} (z(t)) v(t) - (D_+^{\alpha}z(t)) r(t)v(t) \\ &= [-r(t) (D_+^{\alpha}V(t)) - p(t)F (K(t))] v(t) - (D_+^{\alpha}z(t)) r(t)v(t) \quad (21) \\ &< 0, \quad t \geq t_1. \end{aligned}$$

Therefore, $(D_+^{\alpha}z(t)) v(t)$ is a strictly decreasing in $[t_1, \infty)$. We claim that $D_+^{\alpha}z(t) > 0$ for $t \geq t_1$. If not, then $D_+^{\alpha}z(t) \leq 0$ for $t \geq t_1$. Therefore, there exists a $T \geq t_1$ such that

$$(D_+^{\alpha}z(t)) v(t) < (D_+^{\alpha}z(T)) v(T) := -C < 0, \quad t > T. \quad (22)$$

Using Lemma 2.1 in (3.13), we get

$$\frac{K'(t)}{\Gamma(1-\alpha)} = D_+^\alpha z(t) < \frac{-C}{v(t)} = -C \exp\left(\int_{t_0}^t r(s)ds\right)$$

$$\exp\left(\int_{t_0}^t r(s)ds\right) < -\frac{K'(t)}{C\Gamma(1-\alpha)}, \quad t \geq T. \quad (23)$$

Now, we integrating (3.14) from T to t, we have

$$\int_T^t \exp\left(\int_{t_0}^s r(s)ds\right) dt < \frac{-K(t) + K(T)}{C\Gamma(1-\alpha)}$$

$$< \frac{K(T)}{C\Gamma(1-\alpha)} := K_1. \quad (24)$$

Letting $t \rightarrow \infty$, which contradicts to (3.1). Hence $D_+^\alpha z(t) \geq 0$ for $t \geq t_1$ holds. We define the generalized Riccati function $w(t)$ by

$$w(t) = \frac{(D_+^\alpha z(t)v(t))}{F(K(t))}, \quad t \geq t_1. \quad (25)$$

we have $w(t) > 0$ and

$$w'(t) = \frac{((D_+^\alpha z(t)v(t)))'}{F(K(t))} - \frac{(D_+^\alpha z(t))v(t)F'(K(t))K'(t)}{F^2(K(t))}$$

$$\leq \frac{(D_+^{1+\alpha} z(t)v(t) - (D_+^\alpha z(t))v(t)r(t))}{F(K(t))} - \frac{\mu\Gamma(1-\alpha)D_+^\alpha z(t)v(t)D_+^\alpha z(t)}{F^2(K(t))}$$

$$\leq \frac{-r(t)(D_+^\alpha V(t))w(t)}{D_+^\alpha z(t)} - p(t)v(t) - w(t)r(t) - \frac{\mu\Gamma(1-\alpha)w^2(t)}{v(t)}$$

$$\leq -p(t)v(t) - w(t)r(t) - \frac{\mu\Gamma(1-\alpha)w^2(t)}{v(t)} \quad (26)$$

Taking $m=2$,

$$X = \sqrt{\frac{\mu\Gamma(1-\alpha)}{v(t)}}w(t), \quad Y = \frac{r(t)\sqrt{v(t)}}{2\sqrt{\mu\Gamma(1-\alpha)}}. \quad (27)$$

Using Lemma 2.3 and (3.18) in (3.17), we get

$$w'(t) \leq -p(t)v(t) + \frac{(r(t))^2 v(t)}{4\mu\Gamma(1-\alpha)}.$$

Integrating both sides of the above inequality from t_1 to t , we have

$$\begin{aligned} \int_{t_1}^t \left[p(s)v(s) - \frac{(r(s))^2 v(s)}{4\mu\Gamma(1-\alpha)} \right] ds &\leq - \int_{t_1}^t w'(s) ds \\ &= w(t_1) - w(t) \\ &< w(t_1) \end{aligned}$$

Taking the limit supremum of both sides of the above inequality as $t \rightarrow \infty$ we get

$$\limsup_{t \rightarrow \infty} \int_{t_1}^t \left[p(s)v(s) - \frac{(r(s))^2 v(s)}{4\mu\Gamma(1-\alpha)} \right] ds < w(t_1) < \infty$$

which contradicts (3.2). This completes the proof of Theorem 3.1.

Theorem 3.2 Assume that $(A_1) - (A_5)$, (3.1) hold and

$$\int_{t_0}^{\infty} \left[\rho p(s)v(s) - \frac{v(s)(r(s))^2}{4\Gamma(1-\alpha)} \right] ds = \infty, \quad t_0 > 0. \quad (28)$$

Then every solution of the problem (1.1), (1.2) is oscillatory in G .

Proof: Assume that there is a non-oscillatory solution $u(x, t)$ to the problem (1.1), (1.2) which has no zero in $\Omega \times [t_0, \infty)$ for some $t_0 > 0$. Without loss of generality we may suppose that $u(x, t) > 0$, $u(x, t - \tau_j) > 0$ and $u(x, t - \sigma) > 0$ in $\Omega \times [t_1, \infty)$ for $t_1 \geq t_0$, $j = 1, 2, 3, \dots, l$. Proceeding as in the proof of Theorem 3.1, we get (3.11) and $D_+^\alpha z(t) > 0$ for $t \geq t_1$.

Define the function as follows, $w_1(t)$ by

$$w_1(t) = \frac{D_+^\alpha z(t)v(t)}{K(t)}, \quad t \geq t_1. \quad (29)$$

Then, $w_1(t) > 0$,

$$\begin{aligned} w_1'(t) &= \frac{((D_+^\alpha z(t))v(t))'}{K(t)} - \frac{(D_+^\alpha z(t))v(t)K'(t)}{K^2(t)} \\ &= \frac{(D_+^{1+\alpha} z(t))v(t) - (D_+^\alpha z(t))v(t)r(t)}{K(t)} - \frac{\Gamma(1-\alpha)D_+^\alpha z(t)v(t)D_+^\alpha z(t)}{K^2(t)} \\ &\leq \frac{-r(t)(D_+^\alpha V(t))v(t)}{K(t)} - \rho p(t)v(t) - w_1(t)r(t) - \frac{\Gamma(1-\alpha)w_1^2(t)}{v(t)} \end{aligned} \quad (30)$$

$$\begin{aligned} &\leq -\rho p(t)v(t) - w_1(t)r(t) - \frac{\Gamma(1-\alpha)w_1^2(t)}{v(t)} \\ &\leq -\rho p(t)v(t) + \frac{v(t)(r(t))^2}{4\Gamma(1-\alpha)}, \quad t \geq t_1. \end{aligned} \quad (31)$$

By integrating (3.22) from t_1 to t , we have

$$\begin{aligned} \int_{t_1}^t \left[\rho p(s)v(s) - \frac{v(s)(r(s))^2}{4\Gamma(1-\alpha)} \right] ds &\leq - \int_{t_1}^t w_1'(s) ds \\ &= w_1(t_1) - w_1(t) \\ &= w_1(t_1) < \infty, \end{aligned}$$

which contradicts (3.19). Hence the proof is complete. Next, we will discuss some new oscillation criteria for (1.1) by using integral averaging conditions of Philo's type.

Theorem 3.3 Suppose that the conditions $(A_1) - (A_5)$ hold. Furthermore, suppose that there exists a function $H \in C(D, \mathbb{R})$ where $D := \{(t, s) : t \geq s \geq t_0\}$ such that

1. $H(t, t) = 0$ for $t \geq t_0$,

2. $H(t, s) > 0$ for $(t, s) \in D_0$, where $D_0 = \{(t, s) : t > s \geq t_0\}$.

H has a continuous and non positive partial derivative $H'_s(t, s) = \frac{\partial H(t, s)}{\partial s}$ on D_0 with respect to the second variable and satisfies

$$\limsup_{t \rightarrow \infty} \frac{1}{H(t, t_1)} \int_{t_1}^t H(t, s) \left[\rho p(s)v(s) - \frac{v(s)(r(s))^2}{4\Gamma(1-\alpha)} \right] ds = \infty. \quad (32)$$

Then all solutions of (1.1), (1.2) is oscillatory in G .

Proof: Suppose that $u(x, t)$ is a non-oscillatory solution of (1.1), (1.2). Without loss of generality, we may suppose that $u(x, t)$ is an eventually positive solution of (1.1),

(1.2). We proceeding as in the proof of Theorem 3.2, to get that (3.22)

$$w_1'(t) \leq \left[-\rho p(t)v(t) + \frac{v(t)(r(t))^2}{4\Gamma(1-\alpha)} \right]$$

Multiplying the previous inequality by $H(t,s)$ and integrating from t_1 to t for $t \in [t_0, \infty)$, we obtain

$$\int_{t_1}^t H(t,s) \left[\rho p(s)v(s) - \frac{v(s)(r(s))^2}{4\Gamma(1-\alpha)} \right] ds \leq -[H(t,s)w_1(s)]_{t_1}^t + \int_{t_1}^t H_s'(t,s)w_1(s)ds < H(t,t_1)w_1(t_1).$$

Therefore,

$$\frac{1}{H(t,t_1)} \int_{t_1}^t H(t,s) \left[\rho p(s)v(s) - \frac{v(s)(r(s))^2}{4\Gamma(1-\alpha)} \right] ds < w_1(t_1) < \infty$$

which is contradiction to (3.23). The proof is complete.

4. Oscillation of the problem (1.1), (1.3)

In this section we establish some new criteria for the oscillations of the solutions of the equation (1.1),(1.3). For this we need the following.

The smallest eigenvalue β_0 of the Dirichlet problem

$$\begin{aligned} \Delta\omega(x) + \beta\omega(x) &= 0 \quad \text{in } \Omega \\ \omega(x) &= 0 \quad \text{on } \partial\Omega, \end{aligned}$$

is positive and the corresponding eigen function $\phi(x)$ is positive in Ω .

Theorem 4.1 Suppose that the conditions of Theorem 3.1 hold. Then every solution of the problem (1.1),(1.3) is oscillatory in G .

Proof: Assume to the contrary that there is a non-oscillatory solution $u(x,t)$ to the problem (1.1), (1.3) which has no zero in $\Omega \times [t_0, \infty)$ for some $t_0 > 0$. Without loss of generality we may suppose that $u(x,t) > 0$, $u(x,t - \tau_j) > 0$ and $u(x,t - \sigma) > 0$ in $\Omega \times [t_1, \infty)$ for $t_1 \geq t_0$, $j = 1, 2, 3, \dots, l$. Multiplying both sides of (1.1) by $\phi(x)$ and

integrating over the domain Ω , we get

$$\begin{aligned}
 & \int_{\Omega} D_{+,t}^{1+\alpha} [u(x,t) + \lambda(t)u(x,t-\sigma)] \phi(x) dx + r(t) \int_{\Omega} D_{+,t}^{\alpha} u(x,t) \phi(x) dx \\
 &= a(t) \int_{\Omega} \Delta u(x,t) \phi(x) dx + \sum_{j=1}^l a_j(t) \int_{\Omega} \Delta u(x,t-\tau_j) \phi(x) dx \\
 & - \int_{\Omega} p(x,t) F \left(\int_0^t (t-s)^{-\alpha} u(x,s) ds \right) \phi(x) dx + \int_{\Omega} f(x,t) \phi(x) dx, \quad t \geq t_1. \quad (33)
 \end{aligned}$$

Using Green's formula and (1.3) we get,

$$\begin{aligned}
 \int_{\Omega} \Delta u(x,t) \phi(x) dx &= \int_{\Omega} u(x,t) \Delta \phi(x) dx \\
 &= -\beta_0 \int_{\Omega} u(x,t) \phi(x) dx \leq 0, \quad t \geq t_1 \quad (34)
 \end{aligned}$$

and

$$\begin{aligned}
 \int_{\Omega} \Delta u(x,t-\tau_j) \phi(x) dx &= \int_{\Omega} u(x,t-\tau_j) \Delta \phi(x) dx \\
 &= -\beta_0 \int_{\Omega} u(x,t-\tau_j) \phi(x) dx \leq 0, \quad t \geq t_1, \quad j = 1, 2, 3, \dots, l. \quad (35)
 \end{aligned}$$

Let

$$U(t) = \int_{\Omega} u(x,t) \phi(x) dx, \quad t \geq t_1. \quad (36)$$

By using Jensen's inequality and (A_3) , we get

$$\begin{aligned}
 & \int_{\Omega} p(x, t) F \left(\int_0^t (t-s)^{-\alpha} u(x, s) ds \right) \phi(x) dx \\
 & \geq p(t) F \left(\int_{\Omega} \left(\int_0^t (t-s)^{-\alpha} u(x, s) \phi(x) ds \right) dx \right) \\
 & \geq p(t) F \left(\int_0^t (t-s)^{-\alpha} \left(\int_{\Omega} u(x, s) \phi(x) dx \right) ds \right) \\
 & \geq p(t) F (K(t)), \quad t \geq t_1.
 \end{aligned} \tag{37}$$

In view of (4.2) – (4.5), (4.1) yields,

$$D_+^{1+\alpha} [z(t)] + r(t) D_+^{\alpha} U(t) + p(t) F (K(t)) \leq 0, \quad t \geq t_1. \tag{38}$$

The rest of the proof is similar to that of the Theorem 3.1.

The following Theorems can be proved analogously.

Theorem 4.2 Suppose that the conditions of Theorem 3.2 hold. Then every solution of the problem (1.1),(1.3) is oscillatory in G .

Theorem 4.3 Suppose that the conditions of Theorem 3.3 hold. Then every solution of the problem (1.1),(1.3) is oscillatory in G .

5. Example

In this section, we give some examples to illustrate our main theorem in section 3.

Example 5.1 We consider the fractional partial differential equation

$$\begin{aligned}
 & D_{+,t}^{1+\frac{1}{2}} \left[u(x, t) + \frac{1}{9} u(x, t - \frac{\pi}{2}) \right] + \frac{1}{t} D_{+,t}^{\frac{1}{2}} u(x, t) = \frac{8}{9\sqrt{2}} \Delta u(x, t) + \frac{1}{t} \Delta u(x, t - \frac{3\pi}{4}) \\
 & - \frac{1}{\sqrt{2\pi} [\cos tC(x) + \sin tS(x)]} \int_0^t (t-s)^{-\frac{1}{2}} u(x, s) ds - \frac{10}{9\sqrt{2}} \sin x \sin t + 1 \\
 & \text{for } (x, t) \in G, \text{ where } G = (0, \pi) \times [0, \infty),
 \end{aligned} \tag{39}$$

with the boundary condition

$$u(0, t) = u(\pi, t) = 0, \quad t \geq 0. \tag{40}$$

Take $\alpha = \frac{1}{2}, n = 1, m = 1, \lambda(t) = \frac{1}{9}, \sigma = \frac{\pi}{2}, r(t) = \frac{1}{t}, \tau_1 = \frac{3\pi}{4}, a(t) = \frac{8}{9\sqrt{2}}, a_1(t) =$

^{1*}ovsadha@gmail.com, ²kaviakshita@gmail.com

$\frac{1}{t}, \mu = 1, p(x, t) = \frac{1}{\sqrt{2\pi}[\cos tC(x) + \sin tS(x)]}$,
 where $C(x)$ and $S(x)$ are the Fresnel integrals namely,

$$C(x) = \int_0^x \cos\left(\frac{1}{2}\pi t^2\right) dt, S(x) = \int_0^x \sin\left(\frac{1}{2}\pi t^2\right) dt,$$

$F(u) = u$ and $f(x, t) = -\frac{10}{9\sqrt{2}} \sin x \sin t + 1, t > \sin^{-1}\left(\frac{9\sqrt{2}\pi}{20}\right), \mu = 1$.
 But $|C(x)| \leq \pi$ and $|S(x)| \leq \pi$

$$p(t) = \min_{x \in \bar{\Omega}} = \frac{1}{\pi\sqrt{2\pi}[\cos t + \sin t]}.$$

It is clear that conditions $(A_1) - (A_5)$ hold. $\int_{t_0}^{\infty} \exp\left(\int_{t_0}^t \frac{1}{s} ds\right) dt = \infty$ and $v(t) = \frac{t_0}{t}$.
 Consider,

$$\begin{aligned} \int_{t_1}^t \left[v(s)p(s) - \frac{(r(s))^2 v(s)}{4\mu\Gamma(1-\alpha)} \right] ds &= \int_{t_1}^t \left[\frac{t_0}{s} \frac{1}{\pi\sqrt{2\pi}[\cos s + \sin s]} - \frac{1}{s^3} \frac{t_0}{4\Gamma(\frac{1}{2})} \right] ds \\ &> \int_{t_1}^t \left[\frac{t_0}{s} \frac{1}{\pi\sqrt{2\pi}} - \frac{1}{s^3} \frac{t_0}{4\Gamma(\frac{1}{2})} \right] ds \rightarrow \infty \text{ as } t \rightarrow \infty \end{aligned}$$

It is easy to see that all conditions of Theorem 3.1 are satisfied. Hence every solutions of equations (5.1), (5.2) oscillates in $(0, \pi) \times [0, \infty)$. In fact $u(x, t) = \sin x \cos t$ is such a solution.

Example 5.2

$$\begin{aligned} D_{+,t}^{1+\frac{1}{2}} \left[u(x, t) + \frac{1}{t^{\frac{1}{4}}} u(x, t - 2\pi) \right] &+ \frac{1}{t} D_{+,t}^{\frac{1}{2}} u(x, t) \\ &= \left[\frac{3}{4} \frac{\sqrt{2\pi}}{(\Gamma(\frac{1}{4}))^2} t^{-\frac{7}{4}} + \frac{1}{4\sqrt{2}} t^{-\frac{5}{4}} + \frac{t^{-\frac{1}{4}}}{\sqrt{2}} \right] \Delta u(x, t) \\ &+ \left[\frac{1}{\sqrt{2}} + \frac{\sqrt{2\pi}}{(\Gamma(\frac{1}{4}))^2} t^{-\frac{3}{4}} + \frac{t^{-\frac{1}{4}}}{\sqrt{2}} \right] \Delta u(x, t - \frac{\pi}{2}) \\ &- \frac{\sin t}{2\sqrt{\pi}[\sin tC(x) - \cos tS(x)]} \int_0^t (t-s)^{-\frac{1}{2}} u(x, s) ds - \frac{1}{4\sqrt{2}} t^{-\frac{5}{4}} \cos x \sin t \\ &+ \frac{\cos x}{t\sqrt{2}} [\sin t + \cos t] \text{ for } (x, t) \in G, \text{ where } G = (0, \pi) \times [0, \infty), \end{aligned} \quad (41)$$

with the boundary condition

$$u_x(0, t) = u_x(\pi, t) = 0, \quad t \geq 0. \quad (42)$$

Let $\alpha = \frac{1}{2}, n = 1, m = 1, \lambda(t) = \frac{1}{t^{\frac{1}{4}}}, \sigma = 2\pi, r(t) = \frac{1}{t}, \tau_1 = \frac{\pi}{2},$

$$a(t) = \left[\frac{3}{4} \frac{\sqrt{2\pi}}{(\Gamma(\frac{1}{4}))^2} t^{-\frac{7}{4}} + \frac{1}{4\sqrt{2}} t^{-\frac{5}{4}} + \frac{t^{-\frac{1}{4}}}{\sqrt{2}} \right], a_1(t) = \left[\frac{1}{\sqrt{2}} + \frac{\sqrt{2\pi}}{(\Gamma(\frac{1}{4}))^2} t^{-\frac{3}{4}} + \frac{t^{-\frac{1}{4}}}{\sqrt{2}} \right],$$

$$p(x, t) = \frac{\sin t}{2\sqrt{\pi}[\sin tC(x) - \cos tS(x)]}$$

where $C(x)$ and $S(x)$ are the Fresnel integrals namely,

$$C(x) = \int_0^x \cos\left(\frac{1}{2}\pi t^2\right) dt, S(x) = \int_0^x \sin\left(\frac{1}{2}\pi t^2\right) dt,$$

$F(u) = u$ and

$$f(x, t) = -\frac{1}{4\sqrt{2}} t^{-\frac{5}{4}} \cos x \sin t + \frac{\cos x}{t\sqrt{2}} [\sin t + \cos t], \quad \mu = 1.$$

It is easy to see that

$$p_1(t) = \min_{x \in \bar{\Omega}} = \min_{x \in [0, \pi]} = \frac{\sin t}{2\sqrt{\pi} [\sin tC(x) - \cos tS(x)]},$$

$v(t) = \frac{t_0}{t}$. But $|C(x)| \leq \pi$ and $|S(x)| \leq \pi$. Therefore, $p_1(t) = \frac{\sin t}{2\pi\sqrt{\pi}[\sin t - \cos t]}$.

Consider,

$$\begin{aligned} \int_{t_1}^t \left[v(s)p(s) - \frac{(r(s))^2 v(s)}{4\mu\Gamma(1-\alpha)} \right] ds &= \int_{t_1}^t \left[\frac{t_0 \sin s}{s 2\pi\sqrt{\pi} [\sin s - \cos s]} - \frac{1}{s^3} \frac{t_0}{4\Gamma(\frac{1}{2})} \right] ds \\ &> \int_{t_1}^t \left[\frac{t_0}{s 4\pi\sqrt{\pi}} - \frac{1}{s^3} \frac{t_0}{4\Gamma(\frac{1}{2})} \right] ds \rightarrow \infty \text{ as } t \rightarrow \infty \end{aligned}$$

Thus all the conditions of Theorem (3.1) are satisfied. Then every solution of the problem (5.3), (5.4) oscillates in $(0, \pi) \times [0, \infty)$. Infact $u(x, t) = \cos x \sin t$ is one such solution of (5.3), (5.4).

Example 5.3 we consider the fractional partial differential equation

$$\begin{aligned} D_{+,t}^{1+\frac{1}{2}} \left[u(x, t) + \frac{1}{2} u(x, t - \frac{\pi}{2}) \right] + D_{+,t}^{\frac{1}{2}} u(x, t) &= 2\Delta u(x, t) + 3\Delta u(x, t - \pi) \\ - \frac{1}{4t} \left(\int_0^t (t-s)^{-\frac{1}{2}} u(x, s) ds \right)^2 &- \frac{3x}{4\sqrt{\pi}} \frac{1}{t^{\frac{3}{2}}} + \frac{x}{\sqrt{\pi t}} + x^2 \\ \text{for } (x, t) \in G, \text{ where } G &= (0, \pi) \times [0, \infty), \end{aligned} \quad (43)$$

with the boundary condition

$$u(0, t) = u(\pi, t) = 0, \quad t \geq 0. \quad (44)$$

Here $\alpha = \frac{1}{2}$, $n = 1$, $m = 1$, $\lambda(t) = \frac{1}{2}$, $\sigma = \frac{\pi}{2}$, $r(t) = 1$, $\tau_1 = \pi$, $a(t) = 2$, $a_1(t) = 3$, $p(x, t) = \frac{1}{4t}$, $F(u) = u^2$ and $f(x, t) = -\frac{3x}{4\sqrt{\pi}} \frac{1}{t^{\frac{3}{2}}} + \frac{x}{\sqrt{\pi t}} + x^2$, where $t > \frac{3^{\frac{4}{3}}}{4\pi}$, $\mu = 1$.

It is clear that conditions $(A_1) - (A_5)$ hold. $\int_{t_0}^{\infty} \exp(\int_{t_0}^t ds) dt = \infty$ and $v(t) = e^{t_0-t}$. Consider,

$$\int_{t_1}^t \left[v(s)p(s) - \frac{(r(s))^2 v(s)}{4\mu\Gamma(1-\alpha)} \right] ds = \int_{t_1}^t \left[\frac{e^{t_0} e^{-s}}{2\sqrt{t}} - \frac{e^{t_0} e^{-s}}{4\sqrt{\pi}} \right] < \infty$$

It is easy to see that conditions of Theorem 3.1 are not satisfied. Hence every solutions of equations (5.5), (5.6) need not oscillate in $(0, \pi) \times [0, \infty)$. In fact, $u(x, t) = x$ is not a oscillatory solution.

Remark 1: The results obtain in this paper can be considered as generalization of those results in paper [33]. In fact, $\lambda(t) = 0$, $F(u) = u$, $f(x, t) = 0$. Our equation (1.1) reduces to equation (2.1) in [33].

Remark 2: Note that the results by [33] cannot be applied to the equations (5.1), (5.3) and (5.5). So our results improve the results in [33].

References

- [1] Bainov D, Cui BT and E. Minchev, Forced oscillation of hyperbolic equations with deviating arguments, The journal of computational and applied mathematics, 72, 1996, pp. 309-318.
- [2] Cui BT, Oscillation properties for parabolic equations of neutral type, Commentationes mathematicae universitatis carolinae, 33, 1992, pp.581-588.
- [3] Cui BT, Oscillation properties of the solutions of hyperbolic equations with deviating arguments, Demonstration Math., 29, 1996, pp.61-68.
- [4] Cui BT, Yu YH and Lin SZ, Oscillation of solutions of delay hyperbolic differential equations, Acta mathematicae applicatae sinica, 19, 1996, pp.80-88.(in chinese).
- [5] Lalli BS, Yu YH and Cui BT, Oscillation of hyperbolic equations with functional arguments, Applied mathematics and computation, 53, 1993, pp. 97-110.
- [6] Da-Xue Chen, Oscillatory behavior of a class of fractional differential equations with damping, U.P.B.Scientific bulletin, Series A, 75(1), 2013.

- [7] Feng Q and Meng F, Oscillation of solutions to nonlinear forced fractional differential equations, *Electronic journal of differential equations*, 2013, No.169, 2013, pp.1-10.
- [8] Fu XL, Zhuang W, Oscillation of neutral delay parabolic equations, *J. Math. Anal.Appl.*, 191, 1995, pp.473-489.
- [9] Ganesan V and Sathiskumar M, Oscillation theorems for fractional order neutral differential equations, *International journal of math.sci and engg.Appls*, 10, 2016, pp.23-27.
- [10] Grace SR, Agarwal RP, Wong PJY and Zaffer A, On the oscillation of fractional differential equations, *Fractional calculus and applied Analysis*, 15, No.2, 2012, pp 222-231.
- [11] Hale JK, Partial neutral functional-differential equations, (English. English Summary) *Revue roumaine des mathematiques pures et appliquees*, 39, 4, 1994, pp. 339-344.
- [12] Harikrishnan S, Prakash P and Nieto JJ, Forced oscillation of solutions of a nonlinear fractional partial differential equation, *Applied mathematics and computation*, 254, 2015, pp 14-19.
- [13] Jizhong Wang, Fanwei Meng, and Sanyang Liu, Integral average method for oscillation of second order partial differential equations with delays, *Applied mathematics and computation*, 187, 2007, pp. 815-823.
- [14] Mishev DP and Bainov DD, Oscillation of the solutions of parabolic differential equations of neutral type, *Applied mathematics and computation*, 28, 1998, pp. 97-111.
- [15] Prakash P, Harikrishnan S and Nieto JJ and Kim JH, Oscillation of a time fractional partial differential equation, *Electronic journal of qualitative theory of differential equations*, 15, 2014, pp.1-10.
- [16] Raheem A and Maqbul MD, Oscillation criteria for impulsive partial fractional differential equations, *Computer and mathematics with applications*, 73, No.8, 2017, pp.1781-1788.
- [17] Sadhasivam V and Kavitha J, Forced oscillation for a class of fractional parabolic partial differential equations, *Journal of advances in mathematics*, 11, No.6, 2015, pp.5369-5381.
- [18] Sadhasivam V, Kavitha J and Nagajothi N, Oscillation of neutral fractional order partial differential equations with damping term, *International journal of pure and applied mathematics*, 115, No.9, 2017, pp.47-64.

- [19] Wei Nian Li and Weihong Sheng, Oscillation properties for solutions of a kind of Partial fractional differential equations with damping term, *Journal of nonlinear science and applications*, 2016, pp.1600-1608.
- [20] Zahariev AI and Bainov DD, Oscillating properties of the solution of a class of neutral type functional differential equations, *Bulletin of the Australian mathematical society*, 22, 1980, pp.365-372.
- [21] Wei Nian Li, Forced oscillation criteria for a class of fractional Partial differential equations with damping term, *Mathematical Problems in Engineering*, Article ID 410904, 2015, pp.1-6.
- [22] Abbas S, Benchohra M and NGuerekata GM, *Topics in fractional differential equations*, Springer, Newyork, (2012).
- [23] Das S, *Functional fractional calculus for system identification and controls*, Springer, Berlin, (2012).
- [24] Diethelm K, *The analysis of fractional differential equations*, Springer, Berlin, (2010).
- [25] Györi I and Ladas, *Oscillation theory of delay-differential equations*, Clarendon press, Oxford, (1991).
- [26] Hale JK, *Theory of functional differential equations*, Springer-Verlag, Newyork, (1977).
- [27] Hardy GH, Littlewood JE and Polya G, *Inequalities*, Cambridge university press, Cambridge, (1988).
- [28] Kilbas AA, Srivastava HM, and Trujillo JJ, *Theory and applications of fractional differential equations*, Elsevier science B. V., Amsterdam, The Netherlands, 204, (2006).
- [29] Miller KS and Ross B, *An introduction to the fractional calculus and fractional differential equations*, John Wiley and Sons, New York, (1993).
- [30] Podlubny I, *Fractional differential equations*, Academic press, San Diego, Calif, USA, (1999).
- [31] Saker SH, *Oscillation theory of delay differential and difference equations*, VDM Verlag Dr.Muller Aktengesellschaft and Co, USA, (2010).
- [32] Samko SG, Kilbas AA and Marichev OI, *Fractional integrals and derivatives, Theory and applications*, Gordan and Breach Science publishers, Singapore, (1993).

- [33] Varsha Daftardar-Gejji, Fractional calculus theory and applications, Narosa publishing house pvt.Ltd, (2014).
- [34] Vasily Tarasov E, Fractional dynamics, Springer, (2009).
- [35] Wu J, Theory and applications of partial functional differential equations, Applied mathematical sciences, 119, Springer-Verlag, Newyork, (1996).
- [36] N. Yoshida, Oscillation Theory of partial differential equations, World scientific publishing, Singapore, (2008).
- [37] Zhou Y, Basic theory of fractional differential equations, World scientific, Singapore, (2014).