

A Study on Intuitionistic Fuzzy Baire Spaces

Akila S¹ and Mohamed Ali A²

Received: 31 October 2022/ Accepted: 23 November 2022/ Published online: 09 December 2022

©Sacred Heart Research Publications 2017

Abstract

In this paper, the concepts of intuitionistic fuzzy Baire spaces are introduced and characterizations of intuitionistic fuzzy Baire spaces are studied.

Key words: Intuitionistic fuzzy first category, Intuitionistic fuzzy second category, Intuitionistic fuzzy residual set, Intuitionistic fuzzy Baire spaces.

AMS classification: 03E72, 16Y80, 03B52, 20N25.

1 Introduction

The fuzzy concept has invaded almost all branches of mathematics ever since the introduction of fuzzy sets by L.A.Zadeh. The theory of fuzzy topological space was introduced and developed by C.L.Chang and since then various notions in classical topology have been extended to fuzzy topological space. The idea of “intuitionistic fuzzy set” was first published by Atanassov and many works by the same author and his colleagues appeared in the literature. Later, this concept was generalized to “intuitionistic L - fuzzy sets” by Atanassov and Stoeva. The concept of somewhat fuzzy continuous functions and somewhat fuzzy open hereditarily irresolvable by G.Thangaraj and G.Balasubramanian.

In this paper the concepts of intuitionistic fuzzy Baire spaces are introduced and characterizations of intuitionistic fuzzy Baire spaces are studied.

Definition 1.1 Let X be a nonempty fixed set. An intuitionistic fuzzy set (IFS for short) A is an object having the form $A = \{ \langle x, \mu_A(x), \delta_A(x) \rangle : x \in X \}$ where the function $\mu_A : X \rightarrow I$ and $\delta_A : X \rightarrow I$ denote the degree of membership (namely $\mu_A(x)$) and the degree of non membership ($\delta_A(x)$) of each element $x \in X$ to the set

¹Assistant professor, Department of Mathematics, Kuppam Engineering College, Kuppam 517 425, Chittoor District, Andhra Pradesh, India.
Email:agilanambi27@gmail.com

²Associate professor, Department of Mathematics, Islamiah College (Autonomous) Vaniyambadi 635 752, Tirupattur District, Tamil Nadu India.
Email:mohamedalihazim@gmail.com

A , respectively, and $0 \leq \mu_A(x) + \delta_A(x) \leq 1$ for each $x \in X$.

Definition 1.2 Let X be a nonempty set and the intuitionistic fuzzy sets A and B in the form $A = \{\langle x, \mu_A(x), \gamma_A(x) \rangle : x \in X\}$, $B = \{\langle x, \mu_B(x), \gamma_B(x) \rangle : x \in X\}$. Then

- (a) $A \subseteq B$ iff $\mu_A(x) \leq \mu_B(x)$ and $\gamma_A(x) \geq \gamma_B(x)$ for all $x \in X$;
- (b) $A = B$ iff $A \subseteq B$ and $B \subseteq A$;
- (c) $\bar{A} = \{\langle x, \gamma_A(x), \mu_A(x) \rangle : x \in X\}$;
- (d) $A \cap B = \{\langle x, \mu_A(x) \wedge \mu_B(x), \gamma_A(x) \vee \gamma_B(x) \rangle : x \in X\}$;
- (e) $A \cup B = \{\langle x, \mu_A(x) \vee \mu_B(x), \gamma_A(x) \wedge \gamma_B(x) \rangle : x \in X\}$;
- (f) $[] A = \{\langle x, \mu_A(x), 1 - \mu_A(x) \rangle : x \in X\}$
- (g) $\langle \rangle A = \{\langle x, 1 - \gamma_A(x), \gamma_A(x) \rangle : x \in X\}$.

Definition 1.3 $0_\sim = \{\langle x, 0, 1 \rangle : x \in X\}$ and $1_\sim = \{\langle x, 1, 0 \rangle : x \in X\}$.

Definition 1.4 An intuitionistic fuzzy topology (IFT) on a nonempty set X is a family τ of IFSs in X satisfying the following axioms:

- (i) $0_\sim, 1_\sim \in \tau$,
- (ii) $G_1 \cap G_2 \in \tau$ for any $G_1, G_2 \in \tau$,
- (iii) $\cup G_i \in \tau$ for arbitrary family $\{G_i \mid i \in \Lambda\} \subseteq \tau$.

In this case the ordered pair (X, τ) or simply X is called an intuitionistic fuzzy topological space (IFTS) and each IFS in τ is called an intuitionistic fuzzy open set (IFOS). The complement \bar{A} of an IFOS A in X is called an intuitionistic fuzzy closed set (IFCS) in X .

Definition 1.5 Let A be an intuitionistic fuzzy set in an intuitionistic fuzzy topological space X . Then $\text{int}(A) = \bigcup \{G \mid G \text{ is an IFOS in } X \text{ and } G \subseteq A\}$ is called the intuitionistic fuzzy interior of A ; $\text{cl}(A) = \bigcap \{G \mid G \text{ is an IFCS in } X \text{ and } G \supseteq A\}$ is called the intuitionistic fuzzy closure of A .

Definition 1.6 An intuitionistic fuzzy set A in intuitionistic fuzzy topological space (X, T) is called intuitionistic fuzzy dense if there exists no intuitionistic fuzzy closed set B in (X, T) such that $A \subset B \subset 1_\sim$

Proposition 1.7 If A is an intuitionistic fuzzy nowhere dense set in (X, T) , then A is an intuitionistic fuzzy dense set in (X, T) .

Proposition 1.8 Let A be an intuitionistic fuzzy set. If A is an intuitionistic fuzzy closed set in (X, T) with $\text{IFint } A = 0_{\sim}$, then A is an intuitionistic fuzzy nowhere dense set in (X, T) .

2 Intuitionistic Fuzzy Baire Spaces

Definition 2.1 Let (X, T) be an intuitionistic fuzzy topological space. An intuitionistic fuzzy set A in (X, T) is called intuitionistic fuzzy first category if $A = \bigcup_{i=1}^{\infty} B_i$, where B_i 's are intuitionistic fuzzy nowhere dense sets in (X, T) . Any other intuitionistic fuzzy set in (X, T) is said to be of intuitionistic fuzzy second category.

Definition 2.2 An intuitionistic fuzzy topological space (X, T) is called intuitionistic fuzzy first category space if the intuitionistic fuzzy set 1_{\sim} is an intuitionistic fuzzy first category set in (X, T) . That is, $1_{\sim} = \bigcup_{i=1}^{\infty} A_i$ where A_i 's are intuitionistic fuzzy nowhere dense sets in (X, T) . Otherwise (X, T) will be called an intuitionistic fuzzy second category space.

Proposition 2.3 If A be an intuitionistic fuzzy first category set in (X, T) , then $\bar{A} = \bigcap_{i=1}^{\infty} B_i$ where $\text{IFcl}(B_i) = 1_{\sim}$.

Proof:

Let A be an intuitionistic fuzzy first category set in (X, T) . Then $A = \bigcup_{i=1}^{\infty} A_i$, where A_i 's are intuitionistic fuzzy nowhere dense sets in (X, T) . Now $\bar{A} = \bigcup_{i=1}^{\infty} \bar{A}_i = \bigcap_{i=1}^{\infty} (\bar{A}_i)$. Now A_i is an intuitionistic fuzzy nowhere dense set in (X, T) . Then, by Proposition 1.7, we have \bar{A}_i is an intuitionistic fuzzy dense set in (X, T) . Let us put $B_i = \bar{A}_i$. Then $\bar{A} = \bigcap_{i=1}^{\infty} B_i$ where $\text{IFcl}(B_i) = 1_{\sim}$.

Definition 2.4 Let A be an intuitionistic fuzzy first category set in (X, T) . Then \bar{A} is called an intuitionistic fuzzy residual set in (X, T) .

Definition 2.5 Let (X, T) be an intuitionistic fuzzy topological space. Then (X, T) is said to intuitionistic fuzzy Baire space if $\text{IFint}(\bigcup_{i=1}^{\infty} A_i) = 0_{\sim}$, where A_i 's are intuitionistic fuzzy nowhere dense sets in (X, T) .

Proposition 2.6 If $\text{IFint}(\bigcup_{i=1}^{\infty} A_i) = 0_{\sim}$ where $\text{IFint}(A_i) = 0_{\sim}$ and $A_i \in T$, then (X, T) is an intuitionistic fuzzy Baire space.

proof:

Now $A_i \in T$ implies that A_i is an intuitionistic fuzzy closed set in (X, T) . Since $\text{IFint}(A_i) = 0_{\sim}$. By Proposition 1.8, A_i is an intuitionistic fuzzy nowhere dense set in (X, T) . Therefore $\text{IFint}(\bigcup_{i=1}^{\infty} A_i) = 0_{\sim}$. where A_i 's are intuitionistic fuzzy nowhere dense set in (X, T) . Hence (X, T) is an intuitionistic fuzzy Baire space.

Proposition 2.7 IF $\text{IFcl}(\bigcap_{i=1}^{\infty} A_i) = 1_{\sim}$ where A_i 's are intuitionistic fuzzy dense and intuitionistic fuzzy open sets in (X, T) , then (X, T) is an intuitionistic fuzzy Baire Space.

Now $\text{IFcl}(\bigcap_{i=1}^{\infty} A_i) = 1_{\sim}$ implies that $\overline{\text{IFcl}(\bigcap_{i=1}^{\infty} A_i)} = 0_{\sim}$. Then we have $\text{IFint}(\overline{\bigcap_{i=1}^{\infty} A_i}) = 0_{\sim}$. Which implies that $\text{IFint}(\bigcup_{i=1}^{\infty} \overline{A_i}) = 0_{\sim}$. Let $B_i = \overline{A_i}$. Then $\text{IFint}(\bigcup_{i=1}^{\infty} B_i) = 0_{\sim}$. Now $A_i \in T$ implies that $\overline{A_i}$ is an intuitionistic fuzzy closed set in (X, T) and hence B_i is an intuitionistic fuzzy closed and $\text{IFint}(B_i) = \text{IFint}(\overline{A_i}) = \overline{\text{IFcl}(A_i)} = 0_{\sim}$. Hence By Proposition 1.8, B_i is an intuitionistic fuzzy nowhere dense set in (X, T) . Hence $\text{IFint}(\bigcup_{i=1}^{\infty} B_i) = 0_{\sim}$ where B_i 's are intuitionistic fuzzy nowhere dense sets, implies that (X, T) is an intuitionistic fuzzy Baire space.

Proposition 2.8 Let (X, T) be an intuitionistic fuzzy topological space. Then the following are equivalent

- (i) (X, T) is an intuitionistic fuzzy Baire space.
- (ii) $\text{IFint}(A) = 0_{\sim}$, for every intuitionistic fuzzy first category set A in (X, T) .
- (iii) $\text{IFcl}(B) = 1_{\sim}$, for every intuitionistic fuzzy residual set B in (X, T) .

Proof:

(i) \Rightarrow (ii) Let A be an intuitionistic fuzzy first category set in (X, T) . Then $A = (\bigcup_{i=1}^{\infty} A_i)$ where A_i 's are intuitionistic fuzzy nowhere dense sets in (X, T) . Now $\text{IFint}(A) = \text{IFint}(\bigcup_{i=1}^{\infty} A_i) = 0_{\sim}$. Since (X, T) is an intuitionistic fuzzy Baire space. Therefore $\text{IFint}(A) = 0_{\sim}$.

(ii) \Rightarrow (iii) Let B be an intuitionistic fuzzy residual set in (X, T) . Then \overline{B} is an intuitionistic fuzzy first category set in (X, T) . By hypothesis $\text{IFint}(\overline{B}) = 0_{\sim}$ which implies that $\overline{\text{IFcl}(A)} = 0_{\sim}$. Hence $\text{IFcl}(A) = 1_{\sim}$.

(iii) \Rightarrow (i) Let A be an intuitionistic fuzzy first category set in (X, T) . Then $A = (\bigcup_{i=1}^{\infty} A_i)$ where A_i 's are intuitionistic fuzzy nowhere dense sets in (X, T) . Now A

is an intuitionistic fuzzy first category set implies that \bar{A} is an intuitionistic fuzzy residual set in (X, T) . By hypothesis, we have $IFcl(\bar{A}) = 1_{\sim}$, which implies that $IFint(A) = 1_{\sim}$. Hence $IFint(A) = 0_{\sim}$. That is, $IFint(\bigcup_{i=1}^{\infty} A_i) = 0_{\sim}$, where A_i 's are intuitionistic fuzzy nowhere dense sets in (X, T) . Hence (X, T) is an intuitionsitic fuzzy Baire space.

Proposition 2.9 An intuitionistic fuzzy topological space (X, T) is an intuitionistic fuzzy Baire space if and only if $(\bigcup_{i=1}^{\infty} A_i) = 1_{\sim}$, where A_i 's is an intuitionistic fuzzy closed set in (X, T) with $IFint(A_i) = 0_{\sim}$, implies that $IFint(\bigcup_{i=1}^{\infty} A_i) = 0_{\sim}$.

proof:

Let (X, T) be an intuitionistic fuzzy Baire space. Now A_i is an intuitionistic fuzzy closed in (X, T) and $IFint(A_i) = 0_{\sim}$, implies that A_i is an intuitionistic fuzzy nowhere dense set in (X, T) . Now $\bigcup_{i=1}^{\infty} A_i = 1_{\sim}$ implies that 1_{\sim} is an intuitionistic fuzzy first category set in (X, T) . Since (X, T) is an intuitionistic fuzzy Baire space space, by Proposition 2.8, $IFint(1_{\sim}) = 0_{\sim}$. That is, $IFint(\bigcup_{i=1}^{\infty} A_i) = 0_{\sim}$. Conversely suppose that $IFint(\bigcup_{i=1}^{\infty} A_i) = 0_{\sim}$ where A_i . By Proposition 1.8, A_i is an intuitionistic fuzzy nowhere dense set in (X, T) . Hence $IFint(\bigcup_{i=1}^{\infty} A_i) = 0_{\sim}$ implies that (X, T) is an intuitionistic fuzzy Baire space.

Definition 2.10 Let (X, T) and (Y, S) be any two intuitionistic fuzzy topological spaces. A map $f : (X, T) \rightarrow (Y, S)$ is said to be an intuitionistic fuzzy open if the image of every intuitionistic fuzzy open set A in (X, T) is intuitionistic fuzzy open $f(A)$ in (Y, S) .

Proposition 2.11 Let (X, T) and (Y, S) be any two intuitionistic fuzzy topological spaces. If $f : (X, T) \rightarrow (Y, S)$ is an onto intuitionsitic fuzzy contra continuous and intuitionistic fuzzy open then (Y, S) is an intuitionistic fuzzy Baire space.

Proof:

Let A be an intuitionistic fuzzy first category set in (Y, S) . Then $A = (\bigcup_{i=1}^{\infty} A_i)$ where A_i are intuitionistic fuzzy nowhere dense sets in (Y, S) . Suppose $IFint(A) \neq 0_{\sim}$. Then there exists an intuitionistic fuzzy open set $B \neq 0_{\sim}$ in (Y, S) , such that $B \subseteq A$. Then $f^{-1}(B) \subseteq f^{-1}(A) = f^{-1}(\bigcup_{i=1}^{\infty} A_i) = \bigcup_{i=1}^{\infty} f^{-1}(A_i)$. Hence

$$f^{-1}(B) \subseteq \bigcup_{i=1}^{\infty} f^{-1}(IFcl(A_i)) \tag{1}$$

Since f is intuitionistic fuzzy contra continuous and $\text{IFcl}(A_i)$ is an intuitionistic fuzzy closed set in (Y, S) , $f^{-1}(\text{IFcl}(A_i))$ is an intuitionistic fuzzy open in (X, T) . From Eq (1) we have

$$f^{-1}(B) \subseteq \bigcup_{i=1}^{\infty} f^{-1}(\text{IFcl}(A_i)) = \bigcup_{i=1}^{\infty} \text{IFint}(f^{-1}(\text{IFcl}(A_i))) \quad (2)$$

Since f is intuitionistic fuzzy open and onto, $\text{IFint}(f^{-1}(A_i)) \subseteq f^{-1}(\text{IFint}(A_i))$. From Eq (2), we have $f^{-1}(B) \subseteq \bigcup_{i=1}^{\infty} f^{-1}(\text{IFint}(A_i)) \subseteq \bigcup_{i=1}^{\infty} f^{-1}(0_{\sim}) = 0_{\sim}$. Since A_i is an intuitionistic fuzzy nowhere dense. That is, $f^{-1}(B) \subseteq 0_i$ and hence $f^{-1}(B) = 0_{\sim}$ which implies that $B = 0_{\sim}$, which is a contradiction to $B \neq 0_{\sim}$. Hence $\text{IFint}(A) = 0_{\sim}$ where A is an intuitionistic fuzzy first category set in (Y, S) . Hence by Proposition 2.8, (Y, S) is an intuitionistic fuzzy Baire space.

3 Conclusion

In this paper we have presented a Intuitionistic Baire spaces with important definitions and also proved required prepositions. Also we discussed the characterizations of Intuitionistic fuzzy Baire spaces.

References

- [1] Atanassov K, "Intuitionistic Fuzzy Sets", in: V. Sgurev, Ed., VII ITKR's Session, Sofia June 1983 Central Sci. and Techn. Library, Bulg. Academy of Sciences (1984).
- [2] Atanassov K and Stoeva S, "Intuitionistic L-Fuzzy Sets", in: R. Trappl, Ed., Cybernetics and System Research 2,539-540, (1984) .
- [3] Atanassov K, "Intuitionistic Fuzzy Sets", Fuzzy Sets and Systems 20,87-96, (1986)
- [4] Atanassov K, "Review and new results on Intuitionistic Fuzzy Sets", Preprint MFAIS1-88, Sofia,1-8, (1988) .
- [5] Bayhan S and Coker D, "On T_1 and T_2 separation Axioms in Intuitionistic Fuzzy Topological Space", J. Fuzzy Mathematics, 11(3),581-592, (2003) .
- [6] Chang CL, "Fuzzy topological space", J.Math. Anal. Appl., 24,182-190, (1968) .
- [7] Coker D, "An Introduction to Intuitionistic Fuzzy Topological Spaces", Fuzzy Sets and Systems 88,81-89, (1997)

- [8] Coker D, “An introduction to Fuzzy Subspace in Intuitionistic Fuzzy Topological Space”, J. Fuzzy Math., 4 (1996) 749-764.
- [9] Gruenhage G, Lutzer D, “Baire and Volterra Spaces”, Proc. Amer. Soc. 128: 3115-3124, (2000).
- [10] Thangaraj G and Anjalmoose S, On Fuzzy Baire Space, The Journal of Fuzzy Mathematics 21(3),667-676, (2013) .
- [11] Thangaraj G and Anbazhagan C, “Some Remarks on Fuzzy P-Spaces”, Gen. Math. Notes, 26(1),8-16, (2015) .
- [12] Zadeh LA ,Fuzzy sets, Information and Control 8,338-353, (1965) .